

# VUMC Multidisciplinary Surgical Critical Care

---

---

## Gastrointestinal Stress Ulcer Prophylaxis Guideline:

### Background:

Work by Cooke and colleagues ascribed the risk of overt bleeding to be 4.4% and clinically significant bleeding to be 1.5%. The incidence of clinically significant bleeding appears to be dependent on severity of illness and the type of patient population studied. For example, in perioperative cardiac surgery patients the risk is approximately 0.4%. In stroke patients (who were not mechanically ventilated) the risk is 0.1%. There is a strong relationship between duration of mechanical ventilation, duration of intensive care stay, and incidence of ulceration: patients without coagulopathy and mechanical ventilation had an incidence of bleeding of 0.1% in the earlier Cooke study. Again, duration of care and mechanical ventilation represent markers of severity of illness rather than direct causes of ulceration.

Critically ill patients are at risk of GI hemorrhage from primarily gastric or duodenal ulcers. Increased gastric acidity and a decrease in gastric mucosal barrier is believed to be the cause. The longer the gastric pH remains below 4 the greater the risk of hemorrhage. Patients most at risk include critically ill (sepsis, burn, trauma including neuro-trauma) patients requiring >48 hours of mechanical ventilation, patients with a coagulopathy, prior history of GI hemorrhage, organ dysfunction (renal, hepatic, cardiac), or with hypotension/shock. Overall, we know that there is a good relationship between severity of illness (as determined by, for example, Apache II scores) and incidence of ulceration. Moreover, the longer a patient is in ICU, the more likely they are to have a GI bleed. Patients who are likely to have a number of these risk factors, ie EGS patients, (ventilated, hypotensive, coagulopathic), are more likely to have ulceration and bleeding.

As many as 20% of patients may develop clinical GI hemorrhage and if surgery is required mortality can approach 80%.

The most common complication of stress ulcer prophylaxis is pneumonia. The hypothesis is based upon the concept that higher pH relates to overgrowth of gastric microbes and leads to upper tracheal colonization. This concept partnered with microaspiration of intubated patients lying supine may increase the nosocomial pneumonia rate. The ability to reliably maintain a pH <4 will decrease the rate of pneumonia. Several studies comparing the pneumonia rate when comparing sucralfate, antacids and H2 blockade show either improvement or insignificant trends toward decreasing rates with sucralfate. However, due to the current incidence of outpatient ulcer and reflux reduction therapy and the complexity of administering sucralfate, overall benefit appears to be small.

### Purpose:

Standardize the prevention and care of GI hemorrhage

## Indications for Prophylaxis:

### **High Risk Patient:**

- All patients to receive prophylaxis

### **Moderate Risk Patient:**

- Consider prophylaxis

### **Low Risk Patient or Tolerating PO Diet/Full Gastric Enteral Feeds:**

- NO prophylaxis or discontinue prophylaxis

#### **HIGH RISK:**

- Mechanical Ventilation >48 hours
- Coagulopathy
- History of previous GI hemorrhage
- Current outpatient PUD treatment or prophylaxis
- CNS injury (SAH/CVA – hemorrhagic or ischemic)
- Sepsis with or without organ dysfunction
- Vasopressor/inotropic Rx

#### **MODERATE RISK:**

- Chronic NSAID or aspirin use
- High dose prolonged steroid Rx
- ICU stay >10 days

### Nutritional Guideline and Stress Ulcers:

The administration of gastric nutrition reduces, does not eliminate the risk of GI hemorrhage. Any patient predicted to be mechanically ventilated > 48 hours and *without* a contraindication to gastric enteral nutrition, is encouraged to have nasogastric nutrition initiated within 72 hours of admission when a nasoenteric tube is in-situ.

## Prophylaxis Protocols:

### **High Risk Protocol:**

- Start Pepcid 20 mg IV every 12 hours
  - Adjust dose and interval for Ccr <50 mg/dl
  - *UNLESS* patient on PPI for chronic disease (eg. GERD) requiring this medication as outpatient
    - Start PPI
  - If patient placed on TPN; add famotidine to TPN for 24 hours (40 mg) in lieu of IV dosing

- Monitor with routine gastric pH checks (every 6 hours) to keep pH  $\geq 5$ .
  - If pH  $\leq 4$ , increase dose of famotidine.
  - If continues with pH  $\leq 4$ , convert to PPI.

**Patient Who Develops *Significant*\* GI Hemorrhage Receiving Prophylaxis:**

- IV pantoprazole (Protonix®) load then continuous drip
- Consider endoscopic evaluation
- Check gastric pH every 6 hours and adjust dose as needed
- When no evidence of bleeding for 24 hours, convert to intermittent dosing schedule of pantoprazole

*\*Defined as bleeding that requires transfusion, causes hemodynamic changes and/or decrease in Hgb of  $\geq 1$  gram.*

Clinical Management Guidelines (CMG) have been developed for the Multidisciplinary Surgical Critical Care Service in an attempt to standardize and optimize care. They are based on a combination of accepted critical care practice and recent contributions to the medical literature. CMGs are intended to provide guidelines for the management of the majority of patients, and are not proposed as rules, policies or as a substitute for clinical judgment. Deviations from the CMGs are necessary and expected; all exceptions should be documented in the medical record and discussed with the attending physician.

**Authors:**

**Oscar D. Guillamondegui, MD**

Assistant Professor of Surgery

Trauma and Surgical Critical Care  
Vanderbilt University Medical Center

**Robert M. Pousman, DO**

Associate Professor of Surgery and  
Anesthesiology

Medical Director, SICU  
Trauma and Surgical Critical Care  
Vanderbilt University Medical Center

**SICU Medical Director approved:**

---

Addison May, MD, FACS, FCCM  
Revised August 25, 2005

## *Stress Ulcer Prophylaxis*


### **High Risk Factors (Absolute Indications)**

Mechanical Ventilation > 48 hrs  
Coagulopathy  
Hx. GI Bleed

#### Other Indications for Prophylaxis:

SIRS / SEPSIS / MODS  
Cirrhosis / Hepatic Failure  
Renal Failure  
High Dose Steroids  
ISS > 15  
Age > 55

Disease / Injury Requiring Prolong  
Prophylaxis (6 mons.):  
CNS: TBI / SCI  
Burns  
Short Bowel Syndrome



## **Bibliography:**

1. Pruitt BA, Jr., Foley FD, Moncrief JA. Curling's ulcer: a clinical-pathology study of 323 cases. *Ann Surg* 1970; 172(4):523-539.
2. Cook DJ, Fuller HD, Guyatt GH, Marshall JC, Leasa D, Hall R et al. Risk factors for gastrointestinal bleeding in critically ill patients. Canadian Critical Care Trials Group. *N Engl J Med* 1994; 330(6):377-381.
3. Skillman JJ, Bushnell LS, Goldman H, Silen W. Respiratory failure, hypotension, sepsis, and jaundice. A clinical syndrome associated with lethal hemorrhage from acute stress ulceration of the stomach. *Am J Surg* 1969; 117(4):523-530.
4. Cook D, Guyatt G, Marshall J, Leasa D, Fuller H, Hall R et al. A comparison of sucralfate and ranitidine for the prevention of upper gastrointestinal bleeding in patients requiring mechanical ventilation. Canadian Critical Care Trials Group. *N Engl J Med* 1998; 338(12):791-797.
5. Schuster DP, Rowley H, Feinstein S, McGue MK, Zuckerman GR. Prospective evaluation of the risk of upper gastrointestinal bleeding after admission to a medical intensive care unit. *Am J Med* 1984; 76(4):623-630.
6. Shuman RB, Schuster DP, Zuckerman GR. Prophylactic therapy for stress ulcer bleeding: a reappraisal. *Ann Intern Med* 1987; 106(4):562-567.
7. Rosen HR, Vlahakes GJ, Rattner DW. Fulminant peptic ulcer disease in cardiac surgical patients: pathogenesis, prevention, and management. *Crit Care Med* 1992; 20(3):354-359.
8. Wijdicks EF, Fulgham JR, Batts KP. Gastrointestinal bleeding in stroke. *Stroke* 1994; 25(11):2146-2148.
9. Messori A, Trippoli S, Vaiani M, Gorini M, Corrado A. Bleeding and pneumonia in intensive care patients given ranitidine and sucralfate for prevention of stress ulcer: meta-analysis of randomised controlled trials. *BMJ* 2000; 321(7269):1103-1106.
10. Cook DJ, Walter SD, Cook RJ, Griffith LE, Guyatt GH, Leasa D et al. Incidence of and risk factors for ventilator-associated pneumonia in critically ill patients. *Ann Intern Med* 1998; 129(6):433-440.
11. Heiselman DE, Hulisz DT, Fricker R, Bredle DL, Black LD. Randomized comparison of gastric pH control with intermittent and continuous intravenous infusion of famotidine in ICU patients. *Am J Gastroenterol* 1995; 90(2):277-279.
12. Zuckerman, GR, Cort, D, Shuman, RB. Stress Ulcer Syndrome. *J Intensive Care Med*. 1988;3:21-31.
13. Cook, D, Reeve, BK, Guyatt, G, et al. Stress ulcer prophylaxis in critically ill patients-resolving discordant meta-analyses. *JAMA* 1996;275;308-314.